

A Pre-Protostellar Core in L1551

Jonathan J. Swift¹ and William J. Welch

*Department of Astronomy and Radio Astronomy Laboratory, University of California, 601 Campbell Hall,
Berkeley, CA 94720-3411*

James Di Francesco

*National Research Council of Canada, Herzberg Institute of Astrophysics, 5071 West Saanich Road,
Victoria, BC V9E 2E7, Canada*

ABSTRACT

Large field surveys of NH_3 , C_2S , ^{13}CO and C^{18}O in the L1551 dark cloud have revealed a prolate, pre-protostellar molecular core (L1551-MC) in a relatively quiescent region to the northwest of the well-known IRS 5 source. The kinetic temperature is measured to be 9 K, the total mass is $\sim 2 M_\odot$, and the average particle density is $10^4\text{--}10^5 \text{ cm}^{-3}$. L1551-MC is $2'.25 \times 1'.11$ in projection oriented at a position angle of 133° . The turbulent motions are on the order of the sound speed in the medium and contain 4% of the gravitational energy, E_{grav} , of the core. The angular momentum vector is projected along the major axis of L1551-MC corresponding to a rotational energy of $2.5 \times 10^{-3} \sin^{-2} i |E_{\text{grav}}|$. The thermal energy constitutes about a third of $|E_{\text{grav}}|$ and the virial mass is approximately equal to the total mass. L1551-MC is gravitationally bound and in the absence of strong, $\sim 160 \mu\text{G}$, magnetic fields will likely contract on a ~ 0.3 Myr time scale. The line profiles of many molecular species suggest that the cold quiescent interior is surrounded by a dynamic, perhaps infalling envelope which is embedded within the ambient molecular gas of L1551.

Subject headings: ISM: clouds — ISM: individual(L1551) — stars: formation

1. Introduction

Stars play a vital role in the cosmos, enriching the interstellar medium and providing the basic building blocks for the formation of planets and life. Stars with masses $\lesssim 1 M_\odot$ are vastly more numerous than higher mass stars and contain most of the stellar mass in our Galaxy. Much advancement has been made toward the understanding of low mass star formation due in part to the study of nearby star-forming molecular complexes such as Taurus-Auriga. The dense molecular environments most closely associated with the star formation process have been identified (Beichman et al. 1986; Myers 1987) and characterized in large survey data (Jijina, Myers & Adams 1999, and ref-

erences therein). However, the initial conditions under which protostellar collapse occurs remain poorly understood.

Dicarbon monosulphide is a powerful probe of the dense interstellar medium in star-forming regions containing a multitude of emission lines in the millimeter and centimeter wavebands with no hyperfine structure (Saito et al. 1987), a large Zeeman splitting factor (Shinnaga & Yamamoto 2000) and high critical density (Wolkovitch et al. 1997). Observational and theoretical studies of C_2S in the dense interstellar medium suggest that it traces the early evolutionary stages of pre-protostellar cores (Suzuki et al. 1992; Lee et al. 2003) and is otherwise not detectable in star forming regions (Lai & Crutcher 2000).

In 2001, we conducted a C_2S survey of Lynds

¹js@astro.berkeley.edu

1551 (Lynds 1962; Snell 1981) in search of environments in the early stages of star formation. L1551 is an $\sim 80 M_{\odot}$ dark cloud at a distance of 140 pc (Kenyon, Dobrzycka & Hartmann 1994) associated with more than 20 known pre-main sequence stars (see Palla & Stahler 2002, and references therein). The energetics from the embedded young stellar objects are witnessed as molecular outflows (Snell, Loren & Plambeck 1980; Moriarty-Schieven, Butner & Wannier 1995), jets (Mundt et al. 1990) and numerous Herbig-Haro objects (Devine, Reipurth & Bally 1999).

A single emission region appeared in the northwest quadrant of our C₂S survey which we shall refer to as L1551-MC (Lynds 1551 Molecular Core). While L1551-MC is well separated from the known energetics in L1551, there is a low excitation Herbig-Haro object, HH265, which lies at the edge of the projected C₂S emission (Devine, Reipurth & Bally 1999; Garnavich, Noriega-Crespo & Green 1992). No known energetics in L1551 are directed toward this region (see above references and also Figure 1) and the driving source for HH265 remains a mystery.

The many observations at centimeter and millimeter wavelengths used to investigate L1551-MC are described in §2 from which the morphological and physical properties are derived in §3. A discussion of these results is given in §4, and the conclusions, including future study plans, are summarized in §5.

2. Observations and Reductions

Table 1 displays the general parameters of our observations while the text in this section summarizes the relevant observational information not included in the Table.

2.1. BIMA Observations

2.1.1. $\lambda = 9\text{ mm}$

On 2001 September 2 and 3 a $\sim 25' \times 25'$ region centered on L1551 was mapped in C₂S using the BIMA¹ array in its most compact (D) configuration equipped with 30 GHz receivers (Carl-

strom, Joy & Grego 1996). A 7 pointing mosaic ($\sim 10' \times 10'$) centered on the peak emission detected in the D array configuration was then observed in the B and C array configurations in the fall of 2001 and 2002. Only the C and D array data are used (see §3.3.2) which have sufficient $u-v$ coverage to sample all spatial frequencies between $27''$ and $4'.5$.

These observations utilized the highest resolution correlator configuration at BIMA (Welch et al. 1996) which translates to a 0.054 km s^{-1} velocity resolution at the C₂S transition frequency. Phase calibration was ensured with regular scans of the quasar 0449+113 while the flux scale of the observations was determined with scans of Jupiter and 3C84. All data were reduced with the MIRIAD reduction package (Sault, Teuben & Wright 1995) using standard MIRIAD tasks.

2.1.2. $\lambda = 3\text{ mm}$

The C₂S emission region described above (L1551-MC) was then mapped at the transition frequencies of a number of molecular tracers in the 3 mm band. These mosaics all consist of a 7 pointing, Nyquist sampled hexagonal grid centered on $4^{\text{h}}31^{\text{m}}09^{\text{s}}.1, +18^{\circ}12'09''.5$ (J2000) covering a $\sim 2'.5$ region with 100% sensitivity. These tracks were run at the Hat Creek Radio Observatory in the C and D array configurations between 2002 April and June. In addition to the spectral line information, each correlator configuration contained a total of 800 MHz of continuum which was calibrated and preserved.

The $u-v$ coverage was good at all molecular transition frequencies giving well sampled maps for sky angles $15'' \lesssim \theta \lesssim 100''$ except for CS which was only observed in the C array configuration and is more sparsely sampled over this spatial range. The integration time per mosaic pointing totaled ~ 45 minutes for the CS map and ~ 2 hours for all others. System temperatures were variable from ~ 250 to 650 K . All tracks contained 5 minute scans of 0530+135 or 0449+113 at intervals of $\lesssim 25$ minutes for phase calibration and 5 minute integrations on Saturn and W3OH for flux calibration. All data were reduced with the MIRIAD reduction package.

¹The Berkeley-Illinois-Maryland Association array is operated by the University of California, Berkeley, the University of Illinois, and the University of Maryland with support from the National Science Foundation.

2.2. GBT Observations of NH₃

Single pointing spectra and spectral line maps containing the (1,1) and (2,2) inversion transition frequencies of the NH₃ molecule were obtained with the NRAO² 100 m Robert C. Byrd Green Bank Telescope (GBT) in K-band on the nights of 2004 January 6th and 10th. The spectrometer was set to contain 50 MHz of bandwidth centered on a 23.7088144 GHz sky frequency using 9 levels of sampler resolution with 32768 lags in dual polarization mode. Therefore, both the (1,1) and (2,2) inversion transition frequencies including the quadrupole satellites fell within the spectral window with 0.019 km s⁻¹ spectral resolution. The absolute reference position was chosen to lie north of L1551 at 4^h31^m20^s, +18°20′00″ (*J*2000).

The spectral line maps are the result of the On-the-Fly (OTF) mapping technique and cover approximately 5′ × 5′ centered on 4^h31^m08^s.2, +18°13′07″.5 (*J*2000). The GBT main beam efficiency, $\eta_B = 0.81$ and the aperture efficiency, $\eta_A = 0.58$ in K-band. The 60 arcsec s⁻¹ scan rate and 13″ row spacing used to make the maps sufficiently sample all spatial frequencies available to the GBT within the mapped region. System temperature calibrations performed every two rows minimize gain variations across the individual maps and alternating the map scan directions sequentially between right ascension and declination minimize systematic errors (*e.g.* striping). Typical system temperatures were ~ 35 K on both observing nights.

The single pointing spectra were calibrated and exported into ASCII tables with the AIPS++ software package. All further reductions were performed using the IDL software of Research Systems, Inc. The OTF maps were calibrated and gridded in AIPS++ then written to FITS format for further processing in IDL.

Our final spectra, including those in the spectral maps, are the combination of the two independent, calibrated polarization channels fit in off-line frequencies with a third-order polynomial. Our final spectral map is the combination by weighted mean of six individual maps. Systematic effects from the OTF mapping technique are noticeable

in the channels of the final map, although reduced significantly in the velocity integrated map. The total flux in the final spectral map was scaled up by 10% to match the total flux in our individual spectra taken at positions within the mapped region.

2.3. Kitt Peak 12 m Observations at $\lambda = 3$ mm

2.3.1. On-the-Fly ¹³CO and C¹⁸O Maps

A 20′ × 20′ region of L1551 was mapped in ¹³CO(1 → 0) on 2000 November 21 and 22 with the Kitt Peak 12 m Telescope³ using the OTF observing technique. The Millimeter Autocorrelator (MAC) was configured in 2 IF mode with 8192 channels and 150 MHz bandwidth. The main beam, taken to be a 2-dimensional Gaussian with a full width at half-maximum (FWHM) of 55″, comprises 90% of the full beam pattern. The aperture efficiency of the 12m at this observing frequency is 0.5. The 60 arcsec s⁻¹ scan rate and 19″ row spacing of the maps provide spatial over-sampling. The number of rows per reference scan (OFF) and the number of OFFs per total power calibration varied over the course of the observations to keep the system temperature variations below 3% between measurements. System temperatures on the 21st were very good, ~ 180 K, and decent on the 22nd, ~ 275 K.

A comparable OTF map in C¹⁸O(1 → 0) observed on 2002 January 15 and on three consecutive nights starting on 2002 October 21 covered the central 17′ × 17′ of L1551. The backend setup was identical to the ¹³CO observations. System temperatures were good, ~ 250 K for our January observations and variable for the three nights in October.

The raw OTF data were calibrated, converted to visibility data, baseline fit using a first order polynomial, gridded into an image and then written to FITS file format with the AIPS reduction software. Further analyses were performed using IDL.

²The National Radio Astronomy Observatory is a facility of the National Science Foundation, operated under cooperative agreement by Associated Universities, Inc.

³The Kitt Peak 12 m Telescope is operated by the Arizona Radio Observatory at Steward Observatory, University of Arizona, with partial support from the Research Corporation.

2.3.2. CS and N_2H^+ Spectra

Single pointing spectra were taken on the night of 2002 November 30 at selected points within the C_2S emission region at the transition frequencies of both $\text{N}_2\text{H}^+(1 \rightarrow 0)$ and $\text{CS}(2 \rightarrow 1)$. The frequency throw was set to ± 1.0 MHz for CS and ± 3.0 MHz for N_2H^+ . The MAC was configured in 2 IF mode with 12288 usable channels within a 75 MHz band.

Each spectrum was calibrated and folded using UNIPOPS at the observatory and then output into FITS tables. A third order polynomial baseline fit, the combination of independent polarizations, and further analyses were all done using IDL.

3. Results

3.1. Overview

Figure 1 shows our full ^{13}CO Kitt Peak 12 m map with known outflow sources overlaid in the left hand panel. The right hand panel is a closeup of the localized region of C_2S emission L1551 revealed by our BIMA survey. The ^{13}CO is optically thick in this region and has a relative maximum which coincides with the peak of the C_2S emission. HH265 lies at the eastern edge of the projected C_2S .

Figure 2 shows a prolate, dense core seen in NH_3 spatially overlapping with the C_2S emission. There are no compelling young stellar object candidates near L1551-MC in either radio, infrared, or X-ray data. Figure 3 reveals the morphology of the NH_3 emission to more closely resemble the distribution of C^{18}O than the optically thick ^{13}CO .

Figure 4 shows a comparison of many molecular spectra taken at the peak of the C_2S emission, defined as Point A, relative to the central velocity of the NH_3 emission. All lines in Figure 4, excluding C_2S , are from the 12 m observations; BIMA observations of these same transitions did not result in any detections, indicating the overall emission is spatially smooth. The C_2S emission peaks significantly to the red of the reference velocity. There are signs of self-absorption in both the ^{13}CO and CS spectra. The CS line profile in relation to the central velocity of the NH_3 emission is consistent with an infalling envelope of L1551-MC. The N_2H^+ profile is slightly asymmetric, centered somewhat to the red of the NH_3 central velocity,

but is otherwise consistent with the NH_3 emission profile.

3.2. Is L1551-MC Starless?

Jets, winds and outflows from young stellar sources can dramatically affect the interstellar environment from which they are created. Therefore it is important to determine whether or not a stellar point source is present within a molecular core before the molecular line data are interpreted.

The emission at all *IRAS* wavelengths in L1551 is dominated by the L1551-IRS5/L1551NE and HL/XZ Tau regions and shows no evidence of any other sources within L1551. A recent paper by G  lfalk et al. (2004) reports the results from a deep mid-IR search for young stellar objects in L1551 using the Infrared Space Observatory (ISO) and four sources (sources 20, 23, 29 and 32 of their Table 2) lie within a $3'$ radius of L1551-MC. Source 32 is not considered a potential protostar candidate since it was only marginally detected in the $14.3\,\mu\text{m}$ band and has neither an optical nor near-IR counterpart.

The three sources detected at $6.7\,\mu\text{m}$ are plotted on an $(H - K)$ vs. $(K - m_{6.7})$ color-color diagram (Figure 4 in G  lfalk et al. 2004). Sources 23 and 29 are consistent with dust reddened stars with no infrared excess. Source 20 was not detected in the H band. However, this lower limit on m_H places it above the reddening line at a position consistent with being a reddened main-sequence star.

A SIMBAD search reveals one additional radio source within the $3'$ radius of the peak NH_3 emission, GRL11. This 1.4 GHz continuum source is most likely of extragalactic origin (Giovanardi et al. 2000; Snell & Bally 1986). An X-ray source, BFR12, lies within $4''$ of GRL11 and is likely related (Bally, Feigelson & Reipurth 2003). An additional X-ray source from the Chandra survey of L1551, BFR2, is seen to the southwest of L1551-MC and is also most likely a background object.

The combination of all our 3 mm BIMA continuum data precludes the existence of a compact, $< 10''$, source at ~ 100 GHz brighter than 5.3 mJy with 99% confidence. Given these considerations we conclude that L1551-MC is starless. A plot of the G  lfalk et al. (2004), 2MASS and SIMBAD sources can be seen in Figure 2.

3.3. Morphology

3.3.1. Distribution and Orientation of Dense Gas

The dense, inner regions of cores are traced well with NH_3 emission due to a high abundance under these conditions and a tendency to avoid depletion onto grains (Aikawa, Ohashi & Herbst 2003). In this section the primary morphological features of L1551-MC are characterized based on the NH_3 emission in the region. Several single pointing spectra taken at 23 GHz within the larger L1551 cloud reveal that NH_3 emission is not widespread but rather confined to the regions around L1551NE and IRS5 in addition to the region mapped in this paper (see Figure 3). The results from these spectra are summarized in Table 2.

L1551-MC appears roughly elliptical in projection with an axial ratio, $y = 2$. Figure 5 shows the velocity integrated $\text{NH}_3(1,1)$ emission in the optically thin satellite components as contours. This emission distribution is fit with a 2-dimensional Gaussian with the peak emission amplitude, 2-d centroid, major and minor widths, and position angle as free parameters. The gross morphological features, including the characteristic quantities from the average spectrum within a half-maximum NH_3 contour are summarized in Table 3.

A first moment map is created by fitting each spectrum in our NH_3 data cube according to §A. Local velocity gradients calculated at each pixel by vector sum of the central velocity difference at the eight neighboring pixels in the first moment map are shown in Figure 5. There is some scatter to the local gradients, but there is an overall trend of higher velocities toward the southwest. Fitting a velocity gradient to the map in a manner similar to that done by Goodman et al. (1993) yields $\mathcal{G} = 1.2 \pm 0.1 \text{ km s}^{-1} \text{ pc}^{-1}$ at a position angle of $224 \pm 5^\circ$, aligned along the short axis of the velocity integrated emission. The vector addition of all local velocity gradients in Figure 5 yields an identical gradient position angle but smaller amplitude. These results suggest that L1551-MC is prolate. (We assume a prolate spheroidal morphology for L1551-MC in forthcoming calculations unless otherwise stated.)

The velocity integrated emission in all satellite components averaged over elliptical annuli with an

axial ratio of two centered on the (α, δ) coordinates in Table 3 at $13''$ radius intervals is shown in Figure 6. The 2-dimensional projected radial distance from center is defined as b , the impact parameter. The simplest characterization of the intensity profile is of a power law form, $I(b) \propto b^{-p}$. If this function is assumed to describe the intrinsic shape of the intensity profile it becomes immediately apparent that it grossly overestimates the flux in the central pixels. A two component (“broken”) power law function of the form

$$I(b) = \begin{cases} c_1 b^{-p_1} & : b < b_0 \\ c_2 b^{-p_2} & : b > b_0 \end{cases} \quad (1)$$

where $c_2 = c_1 b_0^{(p_2 - p_1)}$, is much more successful in fitting these data. The fit according to the minimum chi-squared value of this intrinsic radial profile convolved with the GBT beam is shown in Figure 6 as the solid line. There is little statistical uncertainty in our best fit parameters, but there is an unknown systematic error since the true 3-dimensional shape of L1551-MC cannot be determined directly from our data. The standard deviation in these parameters invoked by using elliptical annuli of varying axial ratios between $y = 1$ and $y = 2$ gives an estimate of this systematic error. The best fit parameter values are $p_1 = 0.15 \pm 0.04$, $p_2 = 1.48 \pm 0.03$, and $b_0 = 0.040 \pm 0.005 \text{ pc}$.

An intrinsic intensity profile model of the form

$$I(b) = \frac{I_0}{1 + (b/b_0)^\gamma} \quad (2)$$

fits our data to even higher precision than the broken power law. A similar model has been used by Tafalla et al. (2002) to fit the 3-dimensional radial profile of molecular cores with great success and is a good approximation to a Bonner-Ebert profile (Tafalla et al. 2004). This model is used here to fit the 2-dimensional intensity profile only for its convenience and easily interpretable parameters. The minimum chi-squared values for the free parameters of this function are $I_0 = 2.4 \pm 0.2 \text{ K km s}^{-1}$, $b_0 = 0.05 \pm 0.01 \text{ pc}$, and $\gamma = 2.2 \pm 0.1$. This fit is displayed as the dashed line in Figure 6.

A broad brush picture of the mass distribution in L1551-MC can be inferred if it is assumed to be an optically thin prolate spheroid at constant temperature, constant abundance fraction, and in

local thermodynamic equilibrium (LTE). The inferred flat inner density profile with power index ~ 1.2 and break at ~ 0.04 pc where the power law index steepens to ~ 2.5 is typical for a starless core in Taurus (*e.g.* Caselli et al. 2002).

This simple picture is most challenged by the possibilities of a varying NH_3 abundance fraction and a non-symmetric geometry in the third dimension. Recent studies suggest that the abundance fraction of NH_3 increases toward the center of cores (Tafalla et al. 2004), so it is unlikely that an abundance fraction variation alone could produce the flattened inner profile. It would also be unlikely that a geometrical effect could reproduce the effect of having a centrally flat intensity distribution. The existence of a turnover radius at $b_0 \approx 0.04$ pc is therefore robust, while the slope of the profile both inside and outside b_0 is not. No significant N_2H^+ emission is detected in the BIMA 3 mm data, sensitive to spatial scales between 15 and $80''$ at a $3\text{-}\sigma$ level of 0.1 Jy beam^{-1} , suggesting that the densest gas in L1551-MC is distributed smoothly.

3.3.2. The Envelope

Carbon monoxide, the second most abundant molecule in the Galactic interstellar medium, has a relatively small dipole moment and traces low density molecular environments. Figures 1 and 3 show ^{13}CO and C^{18}O to be pervasive in L1551 and therefore these isotopomers probe the low density envelope of L1551-MC as well as the environment in which it sits.

If it is assumed that the rotational excitation temperature of ^{13}CO and C^{18}O are equal, the optical depths of these isotopomers can be estimated. The ratio of antenna temperatures of the ^{13}CO and C^{18}O transitions are

$$\frac{T_A^*(^{13}\text{CO})}{T_A^*(\text{C}^{18}\text{O})} = \frac{(1 - e^{-\tau_{13}}) \eta_{13}}{(1 - e^{-\tau_{18}}) \eta_{18}} \quad (3)$$

where the subscripts 13 and 18 are used to denote quantities related to ^{13}CO and C^{18}O respectively and η is the beam efficiency. The telescope parameters are nearly equivalent at these neighboring transition frequencies so to a good approximation $\eta_{13} = \eta_{18}$. It has been shown that for high column molecular gas ($A_V > 3.5$ mag) the $^{13}\text{CO}/\text{C}^{18}\text{O}$ abundance ratio is relatively uniform and consistent with terrestrial abundance

ratios (McCutcheon et al. 1980) which implies $\tau_{13} = 5.4 \tau_{18}$. Using these numbers, the mean optical depth of ^{13}CO in the NH_3 emission region is 1.9.

The BIMA ^{13}CO and C^{18}O maps show a negligible amount of total emission. There is low level emission visible in the line wings of our ^{13}CO map between the LSR velocities of $5.9\text{--}6.5 \text{ km s}^{-1}$ and $7.2\text{--}7.9 \text{ km s}^{-1}$. However, there is not enough signal above the 0.44 K per channel noise for any interpretation. No emission is detected in the line core where $\tau_{13} > 1$. The peak antenna temperature in our Kitt Peak maps are 5.74 K and 2.23 K in ^{13}CO and C^{18}O respectively and our BIMA maps are well sampled between $15''$ and $80''$ with an RMS of $\sim 0.4 \text{ K}$. Therefore, the ^{13}CO and C^{18}O are tracing a relatively smooth envelope of L1551-MC, with an insignificant amount of structure smaller than $80''$.

The ^{13}CO and C^{18}O emission extends smoothly from the region around L1551-MC into the greater L1551 cloud. The optically thin C^{18}O follows more closely the overall form of NH_3 emission showing no correlation to the C_2S map. Therefore it appears that most of the mass in the envelope follows the gross morphology of the dense inner regions characterized in §3.3.1. This interpretation, however, does not account for depletion of these molecular species onto grains, which is known to occur in dense cores (Aikawa, Ohashi & Herbst 2003; Lee et al. 2003; Tafalla et al. 2004).

The C_2S emission in this region curls around the southwestern limb of the NH_3 emission and stretches to the location of HH265 (see Figure 2). The brightest peak of C_2S emission, defined as Point A in this paper, coincides with a relative maximum in the optically thick ^{13}CO emission (see Figure 1) but shows no such correlation in the optically thin C^{18}O .

No C_2S emission is seen to a $3\text{-}\sigma$ level of $0.27 \text{ Jy beam}^{-1}$ using BIMA in its B array configuration which samples spatial frequencies primarily between $10''$ and $35''$ at the our observing frequency. These data are not used in our final maps and spectra, yet the presented maps reflect an accurate morphology of this tracer given our sensitivity limits.

The CS spectrum at Point A is highly asymmetric containing a blue peak and a red shoulder,

consistent with self-absorption in an infalling layer or shell (Myers et al. 1996; Evans 1999). The C₂S emission is significantly redshifted with respect to the systemic NH₃ velocity. If the C₂S emission region lies on the near side of the core, the arcuate distribution of C₂S may represent an infalling layer around L1551-MC. The possibility of infall in L1551-MC is discussed further in §4.2.

3.4. Physical Properties

Many useful quantities describing conditions within dense molecular environments can be derived from the NH₃(1,1) and (2,2) inversion transitions (see Ho & Townes 1983). Our NH₃(1,1) spectra are fit according to the method described in §A. Minimum chi-squared values less than unity, as expected for a detailed model, were typically achieved using this method and suggest that the LTE assumption is valid for this tracer. Table 4 summarizes the results of our NH₃ fitting and Figure 7 shows fits to the spectra at Point A.

3.4.1. Kinetic Temperature

The flux ratio of the NH₃ (1,1) and (2,2) transitions can be used to estimate the kinetic temperature of the emitting molecular gas (see §A). The best fit parameters for the (1,1) and (2,2) transitions along with Equations A5 and A6 give a kinetic temperature at Point A, $T_k = 8.7 \pm 0.7$ K.

The sensitivity of our NH₃ OTF map was not sufficient to detect the (2,2) transition. While a variation in gas temperature by a factor of ~ 2 has been conjectured for Taurus cores (Galli, Walmsley & Gonçalves 2002), the upper limit on T_k from the Point B spectrum limits the temperature variation to $< 50\%$ (see Table 4). A constant kinetic temperature $T_k = 9 \pm 1$ K within L1551-MC is assumed in all following calculations which include temperature.

3.4.2. Thermal and Turbulent Motions

The full width at half-maximum thermal line width for a gas in LTE at kinetic temperature T_k is given by

$$\Delta v_{th}^2 = 8 \ln 2 \frac{kT_k}{m} \quad (4)$$

where $m = 2.33 m_H$ is the mean particle mass of a gas of molecular hydrogen containing atomic helium with a solar mass fraction ($Y = 0.28$). Using

T_k from §3.4.1, $\Delta v_{th} = 0.42 \text{ km s}^{-1}$. The non-thermal line width of a molecular species can be derived from the observed line width, Δv_{obs} , and kinetic temperature.

$$\Delta v_{nt}^2 = \Delta v_{obs}^2 - 8 \ln 2 \frac{kT_k}{m_{mol}} \quad (5)$$

where m_{mol} is the mass of the observed species. Assuming well mixed conditions, this quantity represents the non-thermal line width for the medium. The total line width can be obtained by adding the thermal and non-thermal components in quadrature.

The mean non-thermal line width is $\Delta v_{nt} = 0.20 \pm 0.002 \text{ km s}^{-1}$ and the mean total line width is $\Delta v_{tot} = 0.46 \pm 0.01 \text{ km s}^{-1}$ obtained by averaging the line widths from each spectrum fit according to §A within a half-maximum NH₃ contour. The errors in the line widths are from Landman, Roussel-Dupré & Tanigawa (1982) and by propagating the error in T_k . The sound speed in the medium is given by, $(kT_k/m)^{1/2} = 0.19 \pm 0.01 \text{ km s}^{-1}$, roughly equal to Δv_{nt} in the region. There is no significant correlation between the projected distance from the peak NH₃ emission and the velocity width.

3.4.3. Mass and Density

The total mass of a molecular core cannot be observed directly and must be inferred using estimates of tracer abundances. The most abundant molecule, H₂, does not radiate in the millimeter or centimeter bands due to its homo-polar geometry, leaving mass estimates of molecular regions dependent on conversion factors for less abundant molecules such as CO. Many processes affect molecular abundance ratios (see van Dishoeck & Blake 1998, for a comprehensive review) and published results for $X(\text{CO})$ in Galactic molecular clouds have varied by a factor of ~ 3 (Lacy et al. 1994). Following is an estimate of the total mass of L1551-MC first using our CO isotopomer data and NH₃ data within a half-maximum contour of the velocity integrated NH₃ intensity and then by consideration of the virial theorem.

The total column of ¹³CO in each pixel of L1551-MC is found assuming that ¹³CO is thermalized at 9 K. These values are corrected for optical depth using Equation 3 and then summed.

Using $X(^{13}\text{CO}) = 1.12 \times 10^{-6}$ converts this number to a total mass of $1.9 M_\odot$.

The total column of NH_3 in the (1,1) state is derived approximating the partition function as the sum over all metastable states up to $J = 6$ (Winnewisser, Churchwell & Walmsley 1979). The total NH_3 mass of L1551-MC is found to be $2.4 \times 10^{-7} M_\odot$ which implies a total molecular mass between 0.8 – $2.4 M_\odot$ for $X(\text{NH}_3)$ between 4.0×10^{-8} (Jijina, Myers & Adams 1999) and 1.5×10^{-8} (Tafalla et al. 2004). The total mass of L1551-MC estimated from molecular line emission is $M_{\text{mol}} \sim 2 M_\odot$ good to within about a factor of 2.

The virial mass is obtained assuming virial equilibrium between thermal and gravitational self-energy. If it is assumed that L1551-MC is comprised of molecular hydrogen

$$M_{\text{vir}} = \frac{25 \Delta v^2 R}{24 \alpha_G G \ln 2} \quad (6)$$

Here α_G is the geometrical and non-uniformity factor of Equation 7 which is of order unity (Bertoldi & McKee 1992), $\Delta v = \Delta v_{\text{tot}}$ from Table 5, and R is the semi-minor axis of the prolate spheroid. Equation 6 yields $M_{\text{vir}} = 1.9^{+0.2}_{-0.7} M_\odot$ for $\alpha_G = 0.87$

An estimate of the volume density from the NH_3 fits is presented in Table 4. This density, n_{ex} , is estimated from the excitation temperature of the fit assuming non-LTE conditions and is a lower limit to the true density of the medium in the derivation (see Equations A7 and following). $\text{N}_2\text{H}^+(1-0)$ is less likely to be in LTE than $\text{NH}_3(1,1)$ owing to a much higher critical density. The hyperfine structure in the $\text{N}_2\text{H}^+(1-0)$ transition is fit in a similar fashion to the NH_3 fits of §A only with the appropriate central frequencies and line strengths for the 7 hyperfine components (Caselli, Myers & Thaddeus 1995; Tiné et al. 2000) input into Equation A4. The excitation temperature from the fit and temperature from §3.4.1 then determine the density of the medium using Equation A7 (see also Caselli et al. 2002). The best fit to the composite N_2H^+ spectrum gives an average particle density $n_{\text{ex}} = 4 \times 10^4 \text{ cm}^{-3}$ in L1551-MC.

The average density inside this core using a mass of $2 M_\odot$ and volume of a prolate spheroid with dimensions from Table 3 is $5 \times 10^5 \text{ cm}^{-3}$. In

consideration of the factor of two error in the mass, the unknown 3-dimensional geometry, and the estimates of n_{ex} from the NH_3 and N_2H^+ data, the average density in L1551-MC $\langle n \rangle \approx 10^4$ – 10^5 cm^{-3} .

4. Discussion

4.1. Energy Partition

The gravitational stability and boundedness of L1551-MC can be estimated through investigation of the energy partition.

The gravitational self-energy for an ellipsoidal mass with a power-law density distribution, $\rho(r) \propto r^{-p}$, is

$$E_{\text{grav}} = -\alpha_G \frac{3}{5} \frac{G M^2}{R} \quad (7)$$

where

$$\alpha_G = \frac{(1 - p/3) \arcsin e}{(1 - 2p/5) e} \quad (8)$$

and e is the eccentricity of the ellipsoid (Bertoldi & McKee 1992). An estimate of the rotational energy can be derived by assuming a spherical geometry with the radius equal to the semi-major axis.

$$E_{\text{rot}} \approx \alpha_I \frac{2}{5} M R_{\text{maj}}^2 \frac{\mathcal{G}^2}{\sin^2 i} \quad (9)$$

where $\mathcal{G}$ is the global velocity gradient and $\alpha_I = (1 - p/3)/(1 - p/5)$. The thermal energy for a diatomic molecular gas is $5/2 N k T_k$ and the turbulent energy is estimated from the non-thermal component of the line width $E_{\text{turb}} = 3/2 M \sigma_{\text{nt}}^2$, where $\sigma_{\text{nt}} = 0.425 \Delta v_{\text{nt}}$.

Table 6 summarizes the results for the energy partition in L1551-MC for three different power law density profiles, $p = 0.0, 1.2$, and 2.0 , using both a spherical, $y = 1$, and prolate spheroidal geometry with $y = 2$. In this table, $E_{\text{vir}} = 2\mathcal{T} + \mathcal{W}$ and represents a measure of the gravitational stability based on virial arguments in the absence of magnetic fields (*e.g.* McKee & Zweibel 1992). In this definition $\mathcal{T} = E_{\text{therm}} + E_{\text{turb}} + E_{\text{rot}}$ is the kinetic energy term in the absence of pressure confinement and $\mathcal{W} = E_{\text{grav}}$ is the gravitational term. The negative values of E_{vir} in Table 6 for all cases is strong evidence that L1551-MC is confined by its own self-gravity. The values most consistent with our analysis in §3 are $p = 1.2$ and $y = 2$.

The strength of the magnetic field is not known in this region. Given the total energy estimates

in Table 6, a field strength of $B \gtrsim 160 \mu\text{G}$ is needed for L1551-MC to be magnetically sub-critical which seems high based on Zeeman measurements (Levin et al. 2001; Crutcher, Heiles & Troland 2003), but is on the order of recent estimates from polarized dust emission in starless cores (Crutcher et al. 2004). The ambient magnetic field position angle in L1551 was measured to be roughly 80° (Vrba, Strom, & Strom 1976) based on the polarization of background stars in the periphery of the cloud. This angle is roughly consistent with the orientation of the jets and outflows and may reflect a relationship between the magnetic field geometry and the stellar formation process in L1551. The minor axis of L1551-MC is also in rough alignment with the ambient magnetic field position angle, but further study of the magnetic field in the immediate vicinity of L1551-MC is needed to determine conclusively its effects on the evolution of this molecular core.

To infer the force balance of a core from energy considerations the equilibrium state must be assumed. From §3.4.3, $M_{\text{mol}} \approx M_{\text{vir}}$ suggesting that L1551-MC is virialized. While virial equilibrium is generally conjectured for ammonia cores (e.g. Jijina, Myers & Adams 1999), recent numerical results suggest that cores are not likely to be hydrostatic and that dynamic cores projected onto the sky can appear in equilibrium (Ballesteros-Paredes, Klessen & Vázquez-Semadeni 2003, and references therein). Additionally, there are no equilibrium conditions for an isolated, self-gravitating, prolate spheroid (Myers et al. 1991). Nevertheless, we can conclude from these analyses that L1551-MC is gravitationally bound and may be unstable. Thermal pressures provide some support against collapse, but ultimately L1551-MC would have to be magnetically sub-critical to prevent collapse on a free-fall time scale, $t_{\text{ff}} \approx 0.3 \text{ Myr}$.

4.2. Core Environment

L1551-MC appears fairly simple traced solely in NH_3 , but is significantly more complex when viewed in other molecular tracers. Figure 4 shows the comparison between the molecular emission of several tracers at Point A with the vertical dotted line signifying the central LSR velocity of the NH_3 (1,1) transition from Table 3. All spectra in Figure 4 are taken in the 3 mm band using the

Kitt Peak 12 m Telescope except the C_2S spectrum which was taken from the BIMA 9 mm map, smoothed to $\sim 55''$ to minimize beam dilution effects.

The ^{13}CO spectrum at the top of Figure 4 is optically thick; $\tau_{13} = 1.9 \pm 0.2$ using Equation 3 with the error determined by Monte Carlo simulation assuming no error in the isotopic ratio. The width of this line is much larger than expected from the direct conversion from the NH_3 width and there is a dip in the line intensity at the central NH_3 velocity. The ^{13}CO is likely tracing a column of gas which extends from the high density interior into the more tenuous and turbulent surroundings.

The CS line in Figure 4 clearly shows a non-Gaussian shape which represents all of our CS spectra taken across the major axis of the C_2S emission. This profile shape has a high peak at $v_{\text{LSR}} \approx 6.5 \text{ km s}^{-1}$ and a shoulder or second peak at $v_{\text{LSR}} \approx 7.0 \text{ km s}^{-1}$ which could be due to another component along the line of sight or self-absorption. There is a $\sim 50\%$ decrement in intensity at the central velocity of the NH_3 emission, suggesting that this feature is due to self-absorption. The broad width of the CS line also means that this gas comprises a larger volume along the line of sight than the NH_3 . It is possible that the asymmetric profile is due to gas infall from a layer (Myers et al. 1996) or spherical shell (Evans 1999).

Another intriguing feature seen in Figure 4 is the redshifted peak of the C_2S emission in relation to the central velocity of the NH_3 . This 0.16 km s^{-1} shift must be due to the kinematics of the gas in L1551-MC since the transition frequency used for our observations has an error (Yamamoto et al. 1990) much less than the observed line width. If the C_2S emission lies on the near side of L1551-MC, this implies that the C_2S is moving toward the highest density environs and may be further evidence of infall. Conversely, this kinematic shift could be due to outflow from L1551-MC. Chemical differentiation within molecular cores is a well known phenomenon (Tafalla et al. 2004, and references therein) and could also be contributing to the discrepancies seen between C_2S and the other molecular tracers. However, there is no clear evidence of depletion in either ^{13}CO or C^{18}O . Alternatively, the C_2S could be tracing a distinct component along the line of

sight. Given the suggestive profile of the CS(2–1) line, the subsonic velocity shift of the C₂S, and the coincidence of the C₂S peak with the optically thick ¹³CO, the interpretation that the C₂S gas is in a state of infall seems most likely.

The N₂H⁺ spectrum in Figure 4 shows the isolated $J_{F1,F} = 1_{0,1} \rightarrow 0_{1,2}$ component at Point A. The fit to the full spectrum (see §3.4.3) yields a value of $\tau_T = 5.5 \pm 0.3$ which means the optical depth of the isolated component is ~ 0.6 . Assuming a beam filling factor of unity, the excitation temperature $T_x = 4.0 \pm 0.1$ K, and the total column of N₂H⁺ within the 12 m beam is $5 \pm 0.5 \times 10^{12} \text{ cm}^{-2}$. The beam corrected column depth ratio of our two nitrogenated molecular tracers, $N(\text{NH}_3)/N(\text{N}_2\text{H}^+) \approx 100$, is consistent with other studies of starless cores in Taurus (Benson, Caselli & Myers 1998). The distribution of N₂H⁺ emission peaks slightly to the red of the NH₃ and is about a factor of 2 broader than that expected from 9 K gas. The N₂H⁺ molecular emission is likely tracing a larger column of gas than the NH₃.

From these spectra, a picture emerges in which the cold, gravitationally bound inner region of L1551-MC is surrounded by a warmer, dynamic envelope showing evidence of gaseous infall. This envelope may not be uniform, but is connected to the more static and smoothly distributed, tenuous molecular gas pervading L1551.

5. Conclusions

In this article, the diagnosis of a newly discovered molecular core found within the L1551 dark cloud is described. This core, referred to as L1551-MC, is located at $4^{\text{h}}31^{\text{m}}09^{\text{s}}.9$, $+18^{\circ}12'41''$ ($J2000$) and has a full width at half-maximum in NH₃ emission of $\sim 2' \times 1'$ at a position angle of 133° east of north. It has an inner region with a relatively flat density profile which turns over to a more steeply declining function at a radius of ~ 0.04 pc. L1551-MC has a mass of $\sim 2 M_{\odot}$, an average density of 10^4 – 10^5 cm^{-3} , and a kinetic temperature of 9 K which varies by less than 50% in the inner region. The velocity of the turbulence is on the order of the sound speed in the medium and comprises 4% of the gravitational energy, E_{grav} , of the core. The rotational energy is $\sim 2.5 \times 10^{-3} \sin^{-2} i |E_{\text{grav}}|$, while the thermal

energy is $0.3 |E_{\text{grav}}|$. The core is gravitationally bound and either will collapse on a time scale of 0.3 Myr or is supported by $\sim 160 \mu\text{G}$ magnetic fields. L1551-MC has a high density interior traced by NH₃ and N₂H⁺ suspended within a dynamic, perhaps infalling envelope which is connected to the more static and tenuous molecular gas seen throughout L1551.

From energy considerations it appears as though L1551-MC will contract to form a star or stellar system. However, there is future work that could further characterize L1551-MC and more accurately place it within the larger star forming context of L1551. An accurate measurement of the magnetic field in L1551-MC will further our understanding of the gravitational support mechanism and constrain the core lifetime. The energy partition depends critically on the total mass and another independent method of determining the total mass, such as from dust emission, should be performed. While the state of the dense, inner regions of L1551-MC are presented herein, supplementary information regarding the envelope of this core is needed. Spectral line maps of CS and its optically thin isotopomer, C³⁴S, will determine the nature of the asymmetric CS line profile and provide a more complete physical, spatial, and chemical picture of the envelope. Lastly, HH265 lies within the projected molecular emission of L1551-MC with no known driving source. Further investigation into the possible relationship between HH265 and L1551-MC may give deeper insight into the evolutionary state of L1551-MC.

We thank Leo Blitz for allocating time for this project during the BIMA 1 cm observing season, and John Carlstrom for donating the usage of his 30 GHz receivers. J. J. S. would like to thank Carl Heiles for thoughtful input. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. This research was partially supported by the National Science Foundation under grant AST 02-28963. Facilities: Berkeley-Illinois-Maryland-Association Array, Green Bank 100 m Telescope, Kitt Peak 12 m Telescope

A. Ammonia Emission Model

We model our observed spectra using

$$T_A^*(\nu) = A(1 - e^{-\tau_\nu}) \quad (\text{A1})$$

where

$$A = \eta_B \eta_f [J_\nu(T_{ex}) - J_\nu(T_{bg})] \quad (\text{A2})$$

and

$$J_\nu(T) = \frac{h\nu}{k} \frac{1}{\exp(h\nu/kT) - 1} \quad (\text{A3})$$

For the (1,1) transition, the optical depth is

$$\tau_\nu = \tau_T \sum_{j=1}^{18} s_j \exp \left[\frac{(\nu - \nu_j)^2}{2\sigma_\nu^2} \right] \quad (\text{A4})$$

where τ_T is the total optical depth of the transition, s_j are the relative line strengths of the 18 hyperfine components, σ_ν is the standard deviation in frequency, and ν_j are the central frequencies of the hyperfine transitions (Kukolich 1967; Rydbeck et al. 1977). To fit the (1,1) transition we wrote a wrapper for the IDL non-linear least-squares fitting routine MPFIT⁴ which easily allows for the tying together of parameters and returns all relevant fitting quantities such as the covariance matrix. More than 70% of the total (2,2) intensity lies in the main component for which there are 3 hyperfine transitions ($\Delta F_1 = 0$ and $\Delta F = 0$) with separations ~ 1 kHz (Kukolich 1967). Therefore, a single Gaussian component is used to fit our (2,2) spectra.

Our non-linear least-squares fitting routine utilizes Equations A1 and A4 to fit our (1,1) spectra with the amplitude factor, A ; the LSR velocity of the emitting gas; the velocity width; and the total optical depth as free parameters. The velocity width is then analytically corrected for optical depth (*e.g.* Phillips et al. 1979) and instrumental broadening. The assumption that the excitation temperatures for all hyperfine transitions are equal is inherent in this fitting procedure.

The relative population of NH_3 molecules in $J = 2$ and $J = 1$ states can be solved for assuming equal excitation temperatures and line widths for the (1,1) and (2,2) transitions (*c.f.* Ho et al. 1979)

$$T_R(2,1) = -T_0 \left[\ln \left\{ \frac{-g_{11}}{g_{22}} \frac{A_{11}}{A_{22}} \left(\frac{\nu_{22}}{\nu_{11}} \right)^2 \frac{1}{\tau_{11} s_{22}} \cdot \ln \left[1 - \frac{T_B(2,2)}{T_B(1,1)} (1 - e^{-\tau_{11} s_{11}}) \right] \right\} \right]^{-1} \quad (\text{A5})$$

where T_0 is the energy difference between the $J = 1$ and $J = 2$ states in units of Kelvin, g_{11} and g_{22} are the statistical weights, τ_{11} is the total optical depth of the (1,1) transition, and s_{11} and s_{22} are the relative line strengths of the transitions.

The kinetic temperature, T_k , can be very well approximated from T_R if $T_k \ll T_0$ since transitions between K-ladders can only happen via collision and non-metastable states within a K-ladder radiatively decay very quickly (Ho & Townes 1983; Walmsley & Ungerechts 1983; Stutzki & Winnewisser 1985). Then, only the (1,1), (2,2) and (2,1) states and their respective radiative and collisional rates need to be considered in the detailed balance equation. Using Ho & Townes (1983) for the spontaneous emission rate and Danby et al. (1988) for the collisional rates, we obtain a transcendental equation for T_k in terms of T_R and T_0

$$T_R = T_k \cdot \left\{ 1 + \frac{T_k}{T_0} \ln [1 + 0.6 \exp(-15.7/T_k)] \right\}^{-1} \quad (\text{A6})$$

⁴<http://astrog.physics.wisc.edu/~craigm/idl/fitting.html>

If a specific value of the beam filling factor is assumed, the excitation temperature can be obtained from the amplitude factor in our fits. The volume particle density can then be estimated using the equation of detailed balance for a 2-level system (Stutzki & Winnewisser 1985; Caselli et al. 2002).

$$\frac{n_{ex}}{n'_{crit}} = \frac{\tilde{T}_x - \tilde{T}_{bg}}{(h\nu/k)(1 - \tilde{T}_x/\tilde{T}_k)} \quad (\text{A7})$$

where

$$n'_{crit} = \beta \cdot \frac{A_{ul}}{C_{ul}}. \quad (\text{A8})$$

A_{ul} and C_{ul} are the spontaneous emission and collisional de-excitation rates for the transition with u and l corresponding to the upper and lower energy levels within the (1,1) state. The escape probability, β should be obtained by detailed modeling, but here we use the approximation,

$$\beta = \frac{1 - e^{-\tau}}{\tau} \quad (\text{A9})$$

where τ is taken to be the maximum optical depth from Equation A4. In Equation A7,

$$\tilde{T} = \left(\frac{h\nu}{k}\right) \left[1 - \exp\left(\frac{-h\nu}{kT}\right)\right]^{-1} \quad (\text{A10})$$

T_k is the kinetic temperature of the medium, $T_{bg} = 2.73 \text{ K}$ is the background radiation temperature, and n_{ex} is the particle density of the collisionally exciting species, taken to be H_2 .

REFERENCES

- Aikawa, Y., Ohashi, N. & Herbst, E. 2003, *ApJ*, 593, 906.
- Caselli, P., Benson, P., Myers, P. & Tafalla, M. 2002, *ApJ*, 572, 238.
- Ballesteros-Paredes, J., Klessen, R. & Vázquez-Semadeni, E. 2003, *ApJ*, 592, 188.
- Bally, J., Feigelson, E. & Reipurth, B. 2003, *ApJ*, 584, 843.
- Beichman, C. A., Myers, P. C., Emerson, J. P., Harris, S., Mathieu, R., Benson, P. J. & Jennings, R. E. 1986, *ApJ*, 307, 337.
- Benson, P., Caselli, P. & Myers, P. 1998, *ApJ*, 596, 743.
- Bertoldi, F. & McKee, C. 1992, *ApJ*, 395, 140.
- Carlstrom, J., Joy, M., Grego, L. 1996, *ApJ*, 456, L75.
- Caselli, P., Myers, P. & Thaddeus, P. 1995, *ApJ*, 455, L77.
- Crutcher, R., Heiles, C. & Troland, T. 2003, in *Lecture Notes in Physics*, vol. 614, ed. Falgarone, E. & Passot, T., Springer, 155.
- Crutcher, R., Nutter, D., Ward-Thompson, D. & Kirk, J. 2004, *ApJ*, 600, 279.
- Danby, G., Flower, D., Valiron, P., Schilke, P. & Walmsley, C. 1988, *MNRAS*, 235, 229.
- Devine, D., Reipurth, B. & Bally, J. 1999, *AJ*, 118, 972.
- Evans, N. 1999, *ARA&A*, 37, 311.
- Gålfalk, M., Oloffson, G., Kaas, A., Olofsson, S., Bontemps, S., Nordh, L. Abergel, A., André, P., Boulanger, F., Burgdorf, M., Casali, M., Cesarsky, C., Davies, J., Falgarone, E., Montmerle, T., Perault, M., Persi, P., Prusti, T., Puget, J. & Sibille, F. 2004, *A&A*, 420, 945.
- Galli, D., Walmsley, M. & Gonçalves, J. 2002, *A&A*, 394, 275.
- Garnavich, P., Noriega-Crespo, A. & Green, P. 1992, *Rev. Mexicana Astron. Astrofis.*, 24, 99.
- Giovanardi, C., Rodríguez, L., Lizano, S. & Cantó, J. 2000, *ApJ*, 538, 728.
- Goodman, A., Benson, P., Fuller, G. & Myers, P. 1993, *ApJ*, 406, 528.
- Ho, P. & Townes, C. 1983, *ARA&A*, 21, 239.
- Ho, P., Barrett, A., Myers, P., Matsakis, D., Cheung, A., Chui, M., Townes, C. & Yngvesson, K. 1979, *ApJ*, 234, 912.
- Jijina, J., Myers, P. & Adams, F. 1999, *ApJS*, 125, 161.
- Kenyon, S., Dobrzycka, D. & Hartmann, L. 1994, *AJ*, 108, 1872.
- Kukolich, S. 1967, *Phys. Rev.*, 156, 83.
- Lacy, J., Knacke, R., Geballe, T. & Tokunaga, A. 1994, *ApJ*, 428, L69.
- Lai, S.-P. & Crutcher, R. 2000, *ApJS*, 128, 271.
- Landman, d., Roussel-Dupré, R. & Tanigawa, G. 1982, *ApJ*, 261, 732.
- Lee, J.-E., Evans, N. J., Shirley, Y. L. & Tatematsu, K. 2003, *ApJ*, 583, 789.
- Levin, S., Langer, W., Velusamy, T., Kuiper, T. & Crutcher, R. 2001, *ApJ*, 555, 850.
- Lynds, B. 1962, *ApJS*, 7, 1.
- McCutcheon, W., Shuter, W., Dickman, R. & Roger, R. 1980, *ApJ*, 237, 9.
- McKee, C. & Zweibel, E. 1992, 399, 551.
- Moriarty-Schieven, G., Butner, H. & Wannier, P. 1995, *ApJ*, 445, L55.
- Mundt, R., Buehrke, T., Solf, J., Ray, T., & Raga, A., 1990, *A&A*, 232, 37.
- Myers, P. C., Fuller, G. A., Mathieu, R. D., Beichman, C. A., Benson, P. J., Schild, R. E. & Emerson, J. P. 1987 *ApJ*, 319, 340.
- Myers, P., Fuller, G., Goodman, A & Benson, P. 1991, *ApJ*, 376, 561.
- Myers, P., Mardones, D., Tafalla, M., Williams, J., Wilner, D. 1996, *ApJ*, 465, L133.

- Phillips, T., Huggins, P., Wannier, P. & Scoville, N. 1979, *ApJ*, 231, 720.
- Pickett, H., Poynter, R., Cohen, E., Delitsky, M., Pearson, J. & Muller, H. 1998, *J. Quant. Spec. Radiat. Transf.*, 60, 883.
- Rydbeck, O., Sume, A., Hjalmarson, Å., Elldér, J., Rönnäng, B. & Kollberg, E. 1977, *ApJ*, 215, L35.
- Saito, S., Kawaguchi, K., Yamamoto, S., Ohishi, M., Suzuki, H. & Kaifu, N. 1987, *ApJ*, 317, 115L.
- Sault, R., Teuben, P. & Wright, M. 1995, in *ASP Conf. Ser. 77, Astronomical Data Analysis Software and Systems IV*, ed., R. A. Shaw, H. E. Payne & J. J. E. Hayes (San Francisco: ASP), 433.
- Shinnaga, H. & Yamamoto, S. 2000, *ApJ*, 544, 330.
- Snell, R., Loren, R. & Plambeck, R., 1980, *ApJ*, 239, 17.
- Snell, R. 1981, *ApJS*, 45, 121.
- Snell, R. & Bally, J. 1986, *ApJ*, 303, 683.
- Palla, F. & Stahler, S. 2002, *ApJ*, 581, 1194.
- Stutzki, J. & Winnewisser, G. 1985, *A&A*, 148, 254.
- Suzuki, H., Yamamoto, S., Ohishi, M., Kaifu, N., Ishikawa, S., Hirahara, Y. & Takano, S. 1992, *ApJ*, 392, 551.
- Tafalla, M., Myers, P., Caselli, P., Walmsley, C. & Comito, C. 2002, *ApJ*, 569, 815.
- Tafalla, M., Myers, P., Caselli & P., Walmsley, C. 2004, *A&A*, 416, 191.
- Tiné, S., Roueff, E., Falgarone, E., Gerin, M. & Pineau des Forêts, G. 2000, *A&A*, 356, 1039.
- van Dishoeck, E. & Blake, G. 1998, *ARA&A*, 36, 317.
- Vrba, F., Strom, S & Strom, K. 1976, *AJ*, 81, 958.
- Walmsley, C. & Ungerechts, H. 1983, *A&A*, 122, 164.
- Welch, W. J. et al. 1996, *PASP*, 108, 93.
- Winnewisser, G., Churchwell, E. & Walmsley, C. 1979, *A&A*, 72, 215.
- Wolkovitch, D., Langer, W., Goldsmith, P. & Heyer, M. 1997, *ApJ*, 477, 241.
- Yamamoto, S., Saito, S., Kawaguchi, K., Chikada, Y., Suzuki, H., Kaifu, N., Ishikawa, S.-I. & Ohishi, M. 1990, *ApJ*, 361, 318.




Fig. 1.— In the right panel, the velocity integrated ^{13}CO in L1551 is overlaid with pre-main sequence stars (stars), an incomplete sample of Herbig-Haro objects (triangles), known jet and outflow directions (arrows) (Mundt et al. 1990; Devine, Reipurth & Bally 1999), and the spatial coverage of our NH_3 map (dashed box). C_2S emission is revealed to the northwest of center in a relatively quiescent region of L1551. In the right hand panel, a close-up view of the C_2S emission region is shown overlaid on the ^{13}CO map which has been restretched to the maximum and minimum values in the region.




Fig. 2.— Velocity integrated NH_3 emission in greyscale with C_2S contours at 30, 50, 70, and 90% the $0.623 \text{ K km s}^{-1}$ maximum overlaid. Boxes signify the point sources from Gálfaik et al. (2004), crosses are points sources from the 2MASS catalog, and the asterisks are SIMBAD objects.




Fig. 3.— Velocity integrated C^{18}O in L1551 with contours of $\text{NH}_3(1,1)$ overlaid at intervals of 30, 50 70 and 90% the peak flux of 2.96 K km s^{-1} . The labeled squares indicate the pointing centers of position switched 23 GHz spectra, the white star symbols are the pre-main sequence stars of Figure 1, and the white triangle is the position of HH265. The Kitt Peak and GBT beams are shown from the bottom left respectively.




Fig. 4.— Spectra from an assortment of molecular transitions at Point A. The CS and N_2H^+ spectra are scaled by $5\times$ and the C_2S spectrum is scaled by $2\times$. The vertical dotted line signifies the systemic velocity of the NH_3 emission at Point A.




Fig. 5.— First moment of NH_3 emission in greyscale is overlaid with local velocity gradients. Arrow lengths signify the relative gradient strengths. The 224° position angle of the global $1.24 \text{ km s}^{-1} \text{ pc}^{-1}$ gradient is shown in the bottom right (length not to scale). Velocity integrated NH_3 intensity of the optically thin satellite components is shown in contour at 30, 50, 70 and 90% maximum intensity.




Fig. 6.— Integrated intensity in the quadrupole satellite components of the $\text{NH}_3(1,1)$ transition as a function of impact parameter. Error bars are the root mean square deviation of the mean intensity within an elliptical annulus and lie mostly within the filled circles. Chi-squared is minimized for models derived from Equations 1 and 2 and the results are plotted as the solid and dashed lines respectively.




Fig. 7.— (a) $\text{NH}_3(1,1)$ spectrum at Point A with §A fit overlaid as the dashed line. The relative line strengths of all 18 hyperfine components are depicted as vertical lines and the LSR velocity scale in each quadrupole window is in reference to the strongest hyperfine line within. (b) The $\text{NH}_3(2,2)$ spectrum of Point A with single Gaussian fit shown as the dashed line. Relative hyperfine line strengths are shown in the main component.

TABLE 1
SUMMARY OF MOLECULAR LINE DATA

BIMA				
Transitions	Frequency (MHz)	$\Delta\nu^a$ (kHz)	Obs. Type	θ_{beam}
C ₂ S(3 ₂ → 2 ₁)	33751.3737 ¹	6.1	Mosaic	27'' × 20''
¹³ CO(1 → 0)	110201.3541 ¹	24.4	Mosaic	10'' × 9''
C ¹⁸ O(1 → 0)	109782.1734 ¹	24.4	Mosaic	10'' × 9''
CS(2 → 1)	97980.9500 ¹	12.2	Mosaic	9'' × 7''
N ₂ H ⁺ (1 → 0)	93176.2650 ²	12.2	Mosaic	11'' × 10''
C ³⁴ S(2 → 1)	109782.1734	12.2	Mosaic	11'' × 10''
Green Bank 100m				
NH ₃ (1,1)	23694.4955 ^{3,4}	3.1	OTF ^b	34''
NH ₃ (1,1)	23694.4955	3.1	APS ^c	34''
NH ₃ (2,2)	23722.6333 ³	3.1	OTF	34''
NH ₃ (2,2)	23722.6333	3.1	APS	34''
Kitt Peak 12m				
¹³ CO(1 → 0)	110201.3541	24.4	OTF	55''
C ¹⁸ O(1 → 0)	109782.1734	24.4	OTF	55''
N ₂ H ⁺ (1 → 0)	93176.2650 ²	12.2	FS ^d	65''
CS(2 → 1)	97980.9500	12.2	FS	65''

^aPost-reduction spectral resolution

^bOn-the-Fly mapping

^cAbsolute Position Switching

^dFrequency Switching

¹From Pickett et al. (1998)

²Frequency of the $J_{F1,F} = 1_{0,1} \rightarrow 0_{1,2}$ hyperfine component (Caselli, Myers & Thaddeus 1995)

³Hyperfine weighted mean transition frequency from Kukolich (1967)

⁴Rydbeck et al. (1977)

TABLE 2
SUMMARY OF THE GBT 23 GHz POSITION SWITCHED OBSERVATIONS

Source	R.A.(J2000)	Dec.(J2000)	$T_B(1,1)^a$	$T_B(2,2)^a$	σ_{T_B}
POINT A	4:31:08.6	+18:12:14.4	1.41	0.23	0.04
POINT B	4:31:04.4	+18:13:30.4	1.09	...	0.07
IRS5	4:31:34.1	+18:08:04.1	2.81	0.53	0.10
L1551NE	4:31:45.1	+18:08:32.0	0.44	...	0.06
XZTAU	4:31:40.0	+18:13:58.0	...	...	0.06
HH30	4:31:37.5	+18:12:26.0	...	...	0.06
OFFXZ	4:31:35.8	+18:14:45.0	...	...	0.06
MIDQNT	4:31:25.0	+18:13:30.0	...	...	0.06
BRTLMB	4:31:15.5	+18:06:55.0	...	...	0.06

NOTE.—Absence of data signifies a non-detection.

^aPeak value of main component brightness temperature in Kelvin.
Beam filling factors are assumed to be unity.

TABLE 3
PROPERTIES OF NH₃ EMISSION IN L1551-MC

R.A. ^a (J2000)	Dec. ^a (J2000)	v_{LSR} ^b (km s ⁻¹)	Δv ^b (km s ⁻¹)	maj. ^c (arcmin)	min. (arcmin)	P.A. degrees
4:31:09.9	+18:12:41	6.72	0.26	2.25	1.11	133

^aCoordinates of the peak emission.

^bThe line profile characteristics are derived from the mean spectrum averaged within a half-maximum contour of the NH₃ emission.

^c“maj.” and “min.” are the FWHM of the major and minor axes respectively.

TABLE 4
NH₃ (1,1) AND (2,2) FIT RESULTS

Source	v_{LSR} (km s ⁻¹)	Δv_{obs} (km s ⁻¹)	T_k^{a} (K)	τ_T	$\eta_f = 1$			$T_x = T_k$	
					T_x (K)	$N(\text{NH}_3)$ (cm ⁻²)	n_{ex}^{b} (cm ⁻³)	$N(\text{NH}_3)$ (cm ⁻²)	η_f
POINT A	6.72	0.24	8.7	5.8	4.4	2.1×10^{15}	5×10^3	3.7×10^{14}	0.26
POINT B	6.71	0.23	< 12.4	2.9	4.5	9.2×10^{14}	$> 4 \times 10^3$	$> 1.5 \times 10^{14}$	> 0.18
IRS5	6.42	0.48	10.0	3.5	6.4	8.2×10^{14}	2×10^4	4.1×10^{14}	0.47
L1551NE	6.55	0.40	< 20.6	4.1	3.2	1.3×10^{16}	$> 8 \times 10^2$	$> 4.6 \times 10^{14}$	> 0.02

^aUpper limits to T_k derived from 3σ level in the (2,2) spectrum. Quantities dependent on T_k were computed with upper limit value when appropriate.

^bDensity derived using equation Equations A7–A9

TABLE 5
L1551-MC PHYSICAL CHARACTERISTICS^a

T_k (K)	M_{mol} ($M_\odot$)	M_{vir} ($M_\odot$)	$\langle n \rangle$ (cm^{-3})	Δv_{th} (km s^{-1})	Δv_{nt} (km s^{-1})	Δv_{tot} (km s^{-1})
9	2	1.9	5×10^4	0.42	0.20	0.46

^aSee text for a discussion of the uncertainties.

TABLE 6
ENERGY PARTITION IN L1551-MC

p^a	y^b	α_G	α_I	E_{grav} (10^{43} ergs)	E_{therm} ($ E_{grav} $)	$E_{turb} \times 10^2$ ($ E_{grav} $)	$E_{rot}^c \times 10^3$ ($ E_{grav} $)	E_{vir}^d ($ E_{grav} $)
0.0	1.0	1.00	1.00	-0.9	0.28	3.2	2.8	-0.37
1.2	1.0	1.15	0.79	-1.1	0.24	2.7	1.9	-0.46
2.0	1.0	1.67	0.56	-1.5	0.17	1.9	0.9	-0.63
0.0	2.0	0.76	1.00	-0.7	0.37	4.1	3.7	-0.18
1.2	2.0	0.87	0.79	-0.8	0.32	3.6	2.5	-0.29
2.0	2.0	1.27	0.56	-1.2	0.22	2.5	1.2	-0.51

^aPower index for $\rho(r) \propto r^{-p}$.

^bAxial ratio, $y = b/a$

^cAn inclination angle, $i = 90^\circ$ is assumed

^d $E_{vir} = 2\mathcal{T} + \mathcal{W}$